

References

- HABU, A., 1981. On Japanese species of *Stomis* (Coleoptera, Carabidae). *Ent. Rev. Japan, Osaka*, **35**: 33–39.
- SCIAKY, R., 1998. Taxonomic review of the genus *Stomis*, with revision of the Chinese species (Coleoptera Carabidae). *Mem. Soc. ent. ital.*, **76**: 21–59.
- STRANEO, S. L., 1953. Nuovi Pterostichini VII (Coleop. Carabidae). *Doriana. Supplemento agli Annli. Mus. civ. Stor. nat. "Giacomo Doria"*, Genova, **1** (36): 1–12.
- TANAKA, K., 1985. Carabidae (Pterostichinae, Zabrinae). In UÉNO, S.-I., Y. KUROSAWA & M. SATÔ (eds.), *Coleoptera of Japan in Color*, **2**: 104–139 [incl. pls. 20–25]. Hoikusha, Osaka. (In Japanese.)

Elytra, Tokyo, **38**(1): 129–130, May 31, 2010

A New Distributional Record of *Habronychus miyatakei* (Coleoptera, Cantharidae)

Naoki TAKAHASHI¹⁾ and Takeo YAMAUCHI²⁾

¹⁾ Entomological Laboratory, Faculty of Agriculture,
Kyushu University, Fukuoka, 812–8581 Japan

²⁾ Toyama Institute of Health, 17–1 Nakataikoyama,
Imizu, Toyama, 939–0363 Japan

Habronychus miyatakei ISHIDA, 1986, was described from Yaku Island, off the southern part of Kyushu. It seems so far endemic to the island, and no collecting record for its distribution has ever been reported from other islands. We will record this species also from Kuchinoerabu Island, lying nearby Yaku Island, for the first time in this short report.

Habronychus (Habronychus) miyatakei ISHIDA, 1986

[Japanese name: Miyatake-higenaga-jôkai]

(Figs. 1–2)

Habronychus miyatakei: M. SATÔ, 1985, 111, pl. 18, fig. 4 (nomen nudum). — ISHIDA, 1986, 218, figs. 11–15 (original description). — TAKAHASHI, 1998, 37 (in list).

Macrohabronychus (Habronychus) miyatakei: BRANCUCCI, 2007, 55 (in list). — KAZANTSEV & BRANCUCCI, 2007, 254 (in list). — N. TAKAHASHI, 2008, 71 (in list).

Specimens examined. 1 ♂, 1 ♀, (Mt.) Shin-dake, Kuchinoerabu Island, Kagoshima Pref., Kyushu, Japan, 13–V–2006, T. YAMAUCHI leg. (preserved in the collection of N. TAKAHASHI).



Figs. 1–2. *Habronychus (Habronychus) miyatakei* ISHIDA, from Kuchinoerabu Is.; 1, male; 2, female.
Scales: 1.0 mm.

Distribution. Japan: Kyushu [Yaku Is., Kuchinoerabu Is. (new record)].

Notes. The generic and subgeneric assignment of this species follows KOPETZ (2008). In comparison with material collected from Yaku Island, the specimens recorded above are somewhat pale in colour, especially on pronotum and scutellum.

We wish to express our gratitude to Dr. Yûichi OKUSHIMA (Kurashiki Museum of Natural History) for his review of the manuscript. This work was supported in part by Global COE Program (Center of excellence for Asian conservation ecology as a basis of human-nature mutualism), MEXT, Japan.

References

- BRANCUCCI, M., 2007. New nomenclatorial and taxonomic acts, and comments: Cantharidae. In LÖBL, I., & A. SMETANA (eds.), *Catalogue of Palaearctic Coleoptera*, 4: 55. Apollo Books, Stenstrup.
- KAZANTSEV, S., & M. BRANCUCCI, 2007. Cantharidae. In LÖBL, I., & A. SMETANA (eds.), *Catalogue of Palaearctic Coleoptera*, 4: 234–298. Apollo Books, Stenstrup.
- KOPETZ, A., 2008. Zur Kenntnis der Familie Cantharidae im Himalaya (Insecta: Coleoptera). *Vernate, Erfurt*, (27): 191–205.
- ISHIDA, K., 1986. The genus *Habronychus* WITTMER of Japan (Coleoptera: Cantharidae). *Trans. Shikoku ent. Soc.*, 17: 215–220.
- SATÔ, M., 1985. Cantharidae. In KUROSAWA, Y., et al. (eds.), *The Coleoptera of Japan in Color*, 3: 107–119 [incl. pls. 17–19]. Hoikusha, Osaka. (In Japanese.)
- TAKAHASHI, K., 1998. Check-list of Cantharidae of Japan. *Kanagawa-Chûhō, Odawara*, (122): 29–48. (In Japanese.)
- TAKAHASHI, N., 2008. A new distributional record of *Malthinellus bicolor* (Coleoptera, Cantharidae, Malthininae) from Yaku Island, Southwest Japan, with a list of cantharids recorded from the island. *Elytra, Tokyo*, 36: 71–72.